

REQUESTED /JK/



United Plaza, Suite 1600
30 South 17th Street
Philadelphia, PA 19103

Telephone: +1-215-568-6400
Facsimile: +1-215-568-6499
www.volpe-koenig.com

BRINGING LAW TO YOUR IDEAS

mail@volpe-koenig.com

FACSIMILE COVER SHEETTO: Examiner Joseph R. Kosack FAX NO.: 571-273-5575FROM: Douglas J. BucklinU.S. Patent Application No. 10/574,982OUR FILE NO. SMC-PT006

Dear Examiner Kosack,

Attached please find correspondence in connection with the above-identified patent application. If you have any questions, please contact us.

NUMBER OF PAGES INCLUDING THIS COVER SHEET: 50

NOTIFY Denise Porreca IF TRANSMISSION IS NOT COMPLETE OR LEGIBLE.

I hereby certify that this paper is being facsimile transmitted to the United States Patent and Trademark Office on February 24, 2010.

A handwritten signature in cursive script that reads 'Denise Porreca'. The signature is written in dark ink and is positioned above the printed name 'Denise Porreca'.

Denise Porreca

2/24/2010
Date

Patents

Trademarks

Copyrights

Trade Secrets

Litigation

Licensing

808741-1

PATENT**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE****In the PATENT APPLICATION of:**

Bansi Lal et al.

Application No.: 10/574,982**Confirmation No.:** 8762**Filed:** April 7, 2006**For:** FIBRINOGEN RECEPTOR
ANTAGONISTS AND THEIR USE**Group:** 1626**Examiner:** Joseph R. Kosack**Our File:** SMC-PT006**Date:** February 24, 2010**PROPOSED EXAMINER'S AMENDMENT**

Examiner Joseph R. Kosack
Group 1626
Facsimile 571-273-5575

Dear Examiner Kosack:

Applicant extends its thanks for considering an Examiner's amendment to place the application in condition for allowance. As per your initial phone call, Applicant is agreeable to cancelling most of the withdrawn claims. But Applicant believes that amendments to claim 32 would allow rejoinder of this claim. Accordingly, Applicant proposes the following amendments and requests that claim 32 be rejoined. As per our subsequent phone call, this facsimile includes a marked claim set showing the proposed amendment, and a clean claim set where all amendments are made. The clean set of claims follows the marked set.


1209407-1

Applicant: Bansi Lal et al.
Application No.: 10/574,982

Marked Claim Set

1 – 24 (Cancelled)


25. (Previously presented) A compound of the general formula (I):



wherein

ring A is phenyl;

R^A is a group of formula (3):



(3)

wherein p is 0;

s is 1;

R¹ is selected from: H, hydroxy, alkyl, partially or fully fluorinated alkyl, alkoxy, alkenyl, alkynyl, carboxy, -C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle;

R³ and R⁴ are independently selected from: H, alkyl, partially or fully fluorinated alkyl, alkenyl, alkynyl, -C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle, -OR⁵, -SR⁵, -NR⁵R⁶, -S(=O)₂NR⁵R⁶, -S(=O)₂R⁵, -C(=O)R⁵, -

Applicant: Bansi Lal et al.
Application No.: 10/574,982

$C(=O)NR^5R^6$, $-C(=O)OR^5$, $-C(=O)SR^5$, $-OC(=O)R^5$, $-OC(=O)OR^5$, $-OC(=O)NR^5R^6$, $-OS(=O)_2R^5$, $-S(C=O)NR^5$ and $-OS(=O)_2NR^5R^6$, or R^3 and R^1 or R^4 , together with the respective nitrogen atoms to which they are attached, form an unsubstituted or substituted 5-, 6- or 7- membered partially saturated or aromatic heterocycle, optionally having one or more additional heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and $-C(=O)OR^5$;

R^5 and R^6 are independently selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein each of said alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkylalkyl group optionally contains at least one hetero atom selected from: N, S and O anywhere in the chain, including the terminal position;

R^B is H;

Y^1 and Y^2 , together, are selected from: $=O$ and $=S$;

Z is N;

W is CH;

R^C is H;

n is 0, 1, 2 or 3;

R^D and R^E are independently selected from: H and an unsubstituted or substituted group selected from: alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkenyl, alkynyl, oxo, carboxy, $-C(=O)OR^5$, $-OR^{17}$, $-SR^{17}$, $-NR^{17}R^{18}$, $-NHC(=O)R^{17}$, $-NHC(=O)OR^{17}$, $-OC(=O)R^{17}$, $-SC(=O)R^{17}$, $-OS(=O)_2R^{17}$ and $-NHS(=O)_2R^{17}$;

R^{17} and R^{18} have the same meaning as R^5 and R^6 , defined above;

R^F is selected from: O, S and $N(OR^{19})$;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

R¹⁹ has the same meaning as R⁵, defined above;

R^G is selected from: aryl, heteroaryl, and partially or fully saturated heterocycle, where said aryl, heteroaryl and heterocycle are substituted by one or more groups of the formula (5):



and optionally, further substituted by one or more groups selected from: -R⁵, halogen, -CN, -SCN, -CNO, -OR²¹, -OC(=O)R²¹, -OS(=O)₂R²¹, -OS(=O)₂NR²¹R²², -OC(=O)OR²¹, -OC(=O)SR²¹, -OC(=O)NR²¹R²², -SR²¹, -S(=O)R²¹, -NO₂, -NR²¹(OR²²), -NR²¹R²², -NR²¹C(=O)R²², -N(R²¹)C(=O)OR²², -N[S(=O)₂R²¹]R²³, C(=O)OR²¹, -S(=O)₂R²¹ and -S(=O)₂OR²¹;

R²¹ has the same meaning as R¹, defined above, and R² is selected from: H, hydroxy, alkyl, partially or fully fluorinated alkyl, alkoxy, alkenyl, alkynyl, carboxy, -C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle:

T is selected from: -CH₂, O, S and NH;

q is 0, 1, 2 or 3;

R²³ and R²⁴ are independently selected from: H, alkyl alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle and C(=O)R²⁵, wherein said alkyl and alkenyl optionally contain at least one hetero atom selected from: O, S and N, in any position of the alkyl or alkenyl chain, and said alkyl and alkenyl are unsubstituted or substituted with at least one group selected from: -OR¹, -OC(=O)R¹, -OS(=O)₂R¹, -S(=O)₂NR¹R², -OC(=O)OR¹, -OC(=O)SR¹, -OC(=O)NR¹R², -SR¹, -S(=O)R¹, -SC(=O)H, -SC(=O)OR¹, -NR¹(OR²), -NR¹R², -NR¹C(=O)R², -N(R¹)C(=O)OR², -NR¹S(=O)₂R², C(=O)OR¹, -S(=O)₂R¹ and -S(=O)₂OR¹;

R²⁵ is selected from: OR⁵, SR⁵, -OCR³R⁴ and -NR⁵R⁶, wherein R³, R⁴, R⁵ and R⁶ are as defined above and wherein optionally, R³ and R⁴, together with the carbon

Applicant: Bansil Lal et al.
Application No.: 10/574,982

to which they are attached, form an unsubstituted or substituted 5-, 6- or 7-membered saturated, partially saturated or aromatic heterocycle having one or more heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and $-C(=O)OR^5$; and the group NR^5R^6 is, optionally, a heterocycle containing at least one additional heteroatom selected from: O, S, and N;

in all its stereoisomeric and tautomeric forms and mixtures thereof in all ratios, and its pharmaceutically acceptable salts.

26. (Previously presented) A compound according to claim 25, wherein R^G is selected from: phenyl, piperidinyl and piperazinyl, and said phenyl, piperidinyl and piperazinyl are substituted by one or more groups of the formula (5):



and optionally, further substituted by one or more groups selected from: $-R^5$, halogen, $-CN$, $-SCN$, $-CNO$, $-OR^{21}$, $-OC(=O)R^{21}$, $-OS(=O)_2R^{21}$, $-OS(=O)_2NR^{21}R^{22}$, $-OC(=O)OR^{21}$, $-OC(=O)SR^{21}$, $-OC(=O)NR^{21}R^{22}$, $-SR^{21}$, $-S(=O)R^{21}$, $-NO_2$, $-NR^{21}(OR^{22})$, $-NR^{21}R^{22}$, $-NR^{21}C(=O)R^{22}$, $-N(R^{21})C(=O)OR^{22}$, $-N[S(=O)_2R^{21}]R^{23}$, $C(=O)OR^{21}$, $-S(=O)_2R^{21}$ and $-S(=O)_2OR^{21}$; and R^{21} and R^{22} are as defined in claim 25.

27. (Previously presented) A compound according to claim 25, wherein R_1 is hydrogen;
 R_3 and R_4 are independently selected from: H, OH, $-C(O)OH$ and $-C(O)Oalkyl$;
 $R^B = R^C = R^D = R^E =$ hydrogen;
 Y^1 and Y^2 , together are $=O$;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

n is the integer 0 or 1;

R^G is phenyl, substituted with one or more of the group of formula (5): T-(CH₂)_q-CR²³R²⁴-COR²⁵, wherein R²³ is H and R²⁴ is H, and, optionally, the compound is further substituted with one or more of the groups selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy, -C(=O)OR⁵, SR²¹, S(=O)₂R²¹ and -N(R²¹)-C(O)CH₃, -CH₂C(O)R²⁵;

and R²⁵ is selected from: OR⁵, OCR³R⁴ and NR⁵R⁶, wherein R³ and R⁴, together with the carbon to which they are attached form an unsubstituted or substituted 5-, 6- or 7- membered saturated, partially saturated or aromatic heterocycle having one or more heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy, -C(=O)OR⁵; and

R⁵, R⁶ and R²¹ are independently selected from: H, alkyl and phenyl.

28. (Currently Amended) A compound according to claim [[1]] 25, wherein

R₁ is hydrogen;

R₃ and R₄ are independently selected from: H, OH, -C(O)OH and -C(O)Oalkyl;

R^B = R^C = R^D = R^E = hydrogen;

Y¹ and Y², together are =O;

n is the integer 0 or 1;

R^G is selected from: piperidinyl and piperazinyl, wherein said piperidinyl and piperazinyl are substituted with one or more of the group of formula (5): T-(CH₂)_q-CR²³R²⁴-COR²⁵, wherein R²³ is H and R²⁴ is H and, optionally, further substituted with one or more groups selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and -C(=O)OR⁵;

and

Applicant: Bansi Lal et al.
Application No.: 10/574,982

R²⁵ is OR⁵, wherein R⁵ is selected from: H, alkyl and phenyl.

29. (Previously presented) A compound according to claim 25 selected from:
- (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
 - (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
 - (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 - (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 - 4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(Benzoyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
 - (4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
 - (4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Benzoyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methanesulfonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-ethoxy carbonyl methoxy-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(imino-{3-methyl-butyrylamino)-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-1-hydroxyimino-ethyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-isobutoxy carbonyl methoxy-phenoxy)-acetic acid isobutyl ester;
2-{4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-NN-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

diethyl-acetamide;

4-(2-[4-(2-(5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl)-acetyl)-phenoxy]-acetoxy)-piperidine-1-carboxylic acid benzyl ester;

4-Benzoyloxycarbonylamino-2-(4-[2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-butyric acid ethyl ester;

4-Benzoyloxycarbonylamino-2-(4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-butyric acid ethyl ester;

(4-[2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenylsulfanyl)-acetic acid methyl ester;

(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-chloro-phenoxy)-acetic acid ethyl ester;

(2-Chloro-4-[2-[5-(imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;

(2-Chloro-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;

(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-ethyl sulfanyl-phenoxy)-acetic acid ethyl ester;

(2-Ethylsulfanyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;

(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-ethane sulfonyl-phenoxy)-acetic acid ethyl ester;

(2-Ethanesulfonyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;

(2,6-Bis-ethylsulfanyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;

(2-Acetyl-amino-4-[2-[5-N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(imino-isobutoxy carbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(N-hydroxy carbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-methoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-propoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy carbonylmethoxy-phenoxy)-acetic acid ethyl ester;
(3-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid;
(2-Ethylsulfonyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Ethyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

2-yl)-acetyl)-phenoxy)-acetic acid ethyl ester;
(5-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-isopropyl-phenoxy)-acetic acid ethyl ester;
(2-*tert*-Butyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Chloro-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Chloro-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-methyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-methyl-phenoxy)-acetic acid benzyl ester;
(2-Ethyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-propyl-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-3-hydroxy-2-propyl-phenoxy)-acetic acid;
(4-Hydroxy-3-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-5-methoxy-phenoxy)-acetic acid ethyl ester;
(3,5-Dihydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-

Applicant: Bansil Lal et al.
Application No.: 10/574,982

yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-3-hydroxy-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-5-hydroxy-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(1-[2S-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-3-(4-hydroxy-phenyl)-propionyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[3-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-propionyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[2-[5-(5-Methyl-isoxazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[2-[5-(*tert*-Butoxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-piperidin-4-yloxy)-acetic acid;
(3-Ethoxy-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(4-[2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-3-ethoxy-phenoxy)-acetic acid ethyl ester;
(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-3-ethoxy-phenoxy)-acetic acid;
(3-Hydroxy-4-[2-[1-oxo-5-(5-oxo-2,5-dihydro-[1,2,4]oxadiazol-3-yl)-1,3-dihydro-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(Acetylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Acetoxy-4-{2-[5-(5-methyl-[1,2,4]oxadiazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid; and
(3-Allyloxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester.

30. (Previously presented) A compound according to claim 27 selected from:
- (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid methyl ester;
 - (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid methyl ester;
 - (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
 - (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
 - (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isopropyl ester;
 - (4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

acetyl)-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-(Benzoyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Benzoyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methanesulfonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-ethoxy carbonyl

Applicant: Bansil Lal et al.
Application No.: 10/574,982

methoxy-phenoxy)-acetic acid ethyl ester;
 (2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 (2-Ethoxycarbonylmethoxy-4-{2-[5-(imino-[3-methyl-butyrylamino]-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 (2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-1-hydroxyimino-ethyl}-phenoxy)-acetic acid ethyl ester;
 (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isobutoxy carbonyl methoxy-phenoxy)-acetic acid isobutyl ester;
 2-(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-NN-diethyl-acetamide;
 4-(2-{4-[2-(5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl)-acetyl]-phenoxy}-acetoxy)-piperidine-1-carboxylic acid benzyl ester;
 4-Benzyloxycarbonylamino-2-(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-butyric acid ethyl ester;
 4-Benzyloxycarbonylamino-2-(4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-butyric acid ethyl ester;
 (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenylsulfanyl)-acetic acid methyl ester;
 (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-chloro-phenoxy)-acetic acid ethyl ester;
 (2-Chloro-4-{2-[5-(imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 (2-Chloro-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
 (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-ethyl sulfanyl-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

phenoxy)-acetic acid ethyl ester;

(2-Ethylsulfanyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-ethanesulfonyl-phenoxy)-acetic acid ethyl ester;

(2-Ethanesulfonyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2,6-Bis-ethylsulfanyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-Acetyl-amino-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(imino-isobutoxy carbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(N-hydroxy carbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;

(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid;

(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

methoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-propoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy carbonylmethoxy-phenoxy)-acetic acid ethyl ester;
(3-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid;
(2-Ethylsulfanyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(5-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isopropyl-phenoxy)-acetic acid ethyl ester;
(2-*tert*-Butyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid benzyl ester;
(2-Ethyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-

Applicant: Bansi Lal et al.
Application No.: 10/574,982

acetyl)-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-2-propyl-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-3-hydroxy-2-propyl-phenoxy)-acetic acid;
(4-Hydroxy-3-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-5-methoxy-phenoxy)-acetic acid ethyl ester;
(3,5-Dihydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(3-Ethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-3-ethoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-3-ethoxy-phenoxy)-acetic acid;
(3-Hydroxy-4-{2-[1-oxo-5-(5-oxo-2,5-dihydro-[1,2,4]oxadiazol-3-yl)-1,3-dihydro-isoindol-2-yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(Acetylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl)-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Acetoxy-4-{2-[5-(5-methyl-[1,2,4]oxadiazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-


Applicant: Bansil Lal et al.
Application No.: 10/574,982

yl]-acetyl)-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid; and
(3-Allyloxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester.

31. (Previously presented) A compound according to claim 28 selected from:
(1-{2S-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-3-(4-hydroxy-phenyl)-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{3-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(5-Methyl-isoxazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(*tert*-Butoxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester; and
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid.

Applicant: Bansilal et al.
Application No.: 10/574,982

32. (Currently amended) A process for the preparation of the compound of claim 25
a compound of having the general formula (I):




wherein all symbols have the same meaning as defined in claim 25.

ring A is phenyl;

R^A is selected from: (CH₂)_pCN, C(=NR¹)SMe and C(=NR¹)OMe, or

R^A is selected from one of the following groups of formula (2), formula (3) and formula (4):



wherein p is 0, 1 or 2;

s is 1 or 2, and when s is 2 the groups R^A are independent of each other and can be the same or different;

R¹ and R² are independently selected from: H, hydroxy, alkyl, partially or fully fluorinated alkyl, alkoxy, alkenyl, alkynyl, carboxy, C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle; or R¹ and R², together with the nitrogen atom to which they are attached, form a saturated, partially saturated or aromatic heterocycle, optionally containing at least one additional hetero atom selected from: N, O and S;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

~~R³ and R⁴ are independently selected from: H, alkyl, partially or fully fluorinated alkyl, alkenyl, alkynyl, C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle, OR⁵, SR⁵, NR⁵R⁶, S(=O)₂NR⁵R⁶, S(=O)₂R⁵, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)SR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, OS(=O)₂R⁵, S(C=O)NR⁵ and OS(=O)₂NR⁵R⁶, or R³ and R¹ or R⁴, together with the respective nitrogen atoms to which they are attached, form an unsubstituted or substituted 5, 6 or 7 membered partially saturated or aromatic heterocycle, optionally having one or more additional heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and C(=O)OR⁵;~~
~~R⁵ and R⁶ are independently selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein each of said alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkylalkyl group optionally contains at least one hetero atom selected from: N, S and O anywhere in the chain, including the terminal position;~~

~~R⁷ and R⁸ have the same meaning as R³ and R⁴, defined above;~~

~~R⁸ is selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein said heterocycle is saturated, partially saturated or aromatic and contains at least one hetero atom selected from: N, O and S, with its point of attachment either through C or N, and wherein each of said alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkylalkyl groups optionally contains at least one hetero atom selected from: N, O and S, anywhere in the chain, including the terminal position;~~

~~R⁹ is selected from: H, halogen, CN, NO₂, alkyl, partially or fully fluorinated alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle, NR¹⁰R¹¹, OR¹⁰, SR¹⁰, S(O)R¹⁰, S(O)₂R¹⁰, NHC(=O)R¹⁰, NHOR¹⁰, OC(=O)R¹⁰,~~

Applicant: Bansil Lal et al.
Application No.: 10/574,982

$\text{SC}(=\text{O})\text{R}^{10}$, $\text{NHC}(=\text{O})\text{OR}^{10}$, $\text{OC}(=\text{O})\text{OR}^{10}$, $\text{C}(=\text{O})\text{NR}^{10}\text{R}^{11}$, $\text{C}(=\text{O})\text{R}^{10}$, and $\text{C}(=\text{O})\text{OR}^{10}$;

R^{10} and R^{11} have the same meaning as R^5 and R^6 , defined above

Y^1 and Y^2 , together, are selected from: $=\text{O}$ and $=\text{S}$;

R^{12} and R^{13} are selected from: H , OR^5 , alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl and aryl;

Z is N ;

W is CH_3 ;

R^C is selected from: H , alkyl, aryl, heterocycle, $=\text{O}$, $=\text{NR}^{14}$, $=\text{S}$, CN , $\text{NR}^{14}\text{R}^{15}$, OR^{14} , SR^{14} , $\text{S}(=\text{O})_2\text{R}^{16}$ and COR^{16} ;

R^{14} and R^{15} have the same meaning as R^5 and R^6 , defined above;

R^{16} is selected from: H , OR^{14} , $\text{N}(\text{R}^{14})_2$, $\text{NR}^{14}\text{R}^{15}$, SR^{14} and R^5 , wherein R^5 , R^{14} and R^{15} are as defined above;

n is 0, 1, 2 or 3;

R^D and R^E are independently selected from: H and an unsubstituted or substituted group selected from: alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkenyl, alkynyl, oxo, carboxy, $\text{C}(=\text{O})\text{OR}^5$, OR^{17} , SR^{17} , $\text{NR}^{17}\text{R}^{18}$, $\text{NHC}(=\text{O})\text{R}^{17}$, $\text{NHC}(=\text{O})\text{OR}^{17}$, $\text{OC}(=\text{O})\text{R}^{17}$, $\text{SC}(=\text{O})\text{R}^{17}$, $\text{OS}(=\text{O})_2\text{R}^{17}$ and $\text{NHS}(=\text{O})_2\text{R}^{17}$;

R^{17} and R^{18} have the same meaning as R^5 and R^6 , defined above;

R^F is selected from: O , S and $\text{N}(\text{OR}^{19})$;

R^{19} and R^{20} have the same meaning as R^5 and R^6 , defined above;

R^G is selected from: aryl, heteroaryl, and partially or fully saturated heterocycle, where said aryl, heteroaryl and heterocycle are substituted by one or more groups of the formula (5):

Applicant: Bansil Lal et al.
Application No.: 10/574,982



and optionally, further substituted by one or more groups selected from: R^5 , halogen, CN, SCN, CNO, OR^{21} , $OC(=O)R^{21}$, $OS(=O)_2R^{21}$, $OS(=O)_2NR^{21}R^{22}$, $OC(=O)OR^{21}$, $OC(=O)SR^{21}$, $OC(=O)NR^{21}R^{22}$, SR^{21} , $S(=O)R^{21}$, $SC(=O)H$, $SC(=O)OR^{21}$, NO_2 , $NR^{21}(OR^{22})$, $NR^{21}R^{22}$, $NR^{21}C(=O)R^{22}$, $N(R^{21})C(=O)OR^{22}$, $NIS(=O)_2R^{21}R^{23}$, $C(=O)OR^{21}$, $S(=O)_2R^{21}$ and $S(=O)_2OR^{21}$; R^{21} and R^{22} have the same meaning as R^1 and R^2 , defined above;

T is selected from: CH_2 , O, S and NH;


q is 0, 1, 2 or 3;

R^{23} and R^{24} are independently selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle and $C(=O)R^{25}$, wherein said alkyl and alkenyl optionally contain at least one hetero atom selected from: O, S and N, in any position of the alkyl or alkenyl chain, and said alkyl and alkenyl are unsubstituted or substituted with at least one group selected from: OR^1 , $OC(=O)R^1$, $OS(=O)_2R^1$, $S(=O)_2NR^1R^2$, $OC(=O)OR^1$, $OC(=O)SR^1$, $OC(=O)NR^1R^2$, SR^1 , $S(=O)R^1$, $SC(=O)H$, $SC(=O)OR^1$, $NR^1(OR^2)$, NR^1R^2 , $NR^1C(=O)R^2$, $N(R^1)C(=O)OR^2$, $NR^1S(=O)_2R^2$, $C(=O)OR^1$, $S(=O)_2R^1$ and $S(=O)_2OR^1$;

R^{25} is selected from: OR^5 , SR^5 , OCR^3R^4 and NR^5R^6 , wherein R^3 , R^4 , R^5 and R^6 are as defined above and wherein optionally, R^3 and R^4 , together with the carbon to which they are attached, form an unsubstituted or substituted 5-, 6- or 7-membered saturated, partially-saturated or aromatic heterocycle having one or more heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and $C(=O)OR^5$; and the group NR^5R^6 is, optionally, a heterocycle containing at least one additional heteroatom selected from: O, S, and N;


Applicant: Bansi Lal et al.
Application No.: 10/574,982

which process comprises
the process comprising: (a) reacting a compound of formula (II):



wherein


L is a leaving group; and all other symbols are as defined above in claim 25; with a compound of the formula (III):



wherein all symbols are as defined above in claim 25;

in the presence of an organic or inorganic base in an organic solvent or a mixture of at least two different organic solvents, at a temperature ranging from -40°C to 150°C , for 0.5 to 16 h, to effect in situ cyclization to form a compound of the general formula (I) above, and, optionally, converting the compound into a physiologically tolerable salt; or


b) reacting a compound of the formula (IV)



Applicant: Bansi Lal et al.
Application No.: 10/574,982


wherein

L_2 is a leaving group; and all other symbols are as defined ~~above~~ in claim 25;
 with a compound of the formula (V):



where R^G is selected from: piperidinyl, piperazinyl and phenyl, wherein said piperidinyl, piperazinyl and phenyl, are optionally substituted with 1, 2, 3 or 4 hydroxyl groups, and all other symbols are as defined ~~above~~ in claim 25, in the presence of an organic or inorganic base in an organic solvent or water at a temperature ranging from 0°C to 150°C, for 0.5 to 12 h, to form a compound of the general formula (I), and, optionally, converting one or more of the hydroxyl groups into a group selected from the substituents for R^G as defined in general formula (I) and, optionally, converting the compound into a physiologically tolerable salt; alternatively, activating a compound of the formula (IV) above, wherein L_2 is -OH, by treatment with a mixed anhydride to form a peptide coupling with a compound of the formula (V), wherein R^G is piperidinyl or piperazinyl, and thereby provide a compound of the general formula (I), wherein R^G is piperidinyl or piperazinyl substituted with at least a group of the formula (5); and, optionally, converting the resultant compound into a physiologically tolerable salt; ~~or~~


~~e) alkylating a compound of the formula (VIII):~~



Applicant: Bansi Lal et al.
Application No.: 10/574,982

wherein B is halogen, acetate or formate, and all other symbols are as defined above;

with a compound of the formula:



wherein

R^G is phenyl, having at least one substituent which is $\text{OCH}_2\text{Phenyl}$, and optionally at least one further substituent selected from: R^5 , halogen, CN, SCN , CNO , OR^{21} , $\text{OC}(=\text{O})\text{R}^{21}$, $\text{OS}(=\text{O})_2\text{R}^{21}$, $\text{OS}(=\text{O})_2\text{NR}^{21}\text{R}^{22}$, $\text{OC}(=\text{O})\text{OR}^{21}$, $\text{OC}(=\text{O})\text{SR}^{21}$, $\text{OC}(=\text{O})\text{NR}^{21}\text{R}^{22}$, SR^{21} , $\text{S}(=\text{O})\text{R}^{21}$, $\text{SC}(=\text{O})\text{H}$, $\text{SC}(=\text{O})\text{OR}^{21}$, NO_2 , NR^{21}OH , $\text{NR}^{21}(\text{OR}^{22})$, $\text{NR}^{21}\text{R}^{22}$, $\text{NR}^{21}\text{C}(=\text{O})\text{R}^{22}$, $\text{N}(\text{R}^{21})\text{C}(=\text{O})\text{OR}^{22}$, $\text{N}[\text{S}(=\text{O})_2\text{R}^{21}]\text{R}^{22}$, $\text{C}(=\text{O})\text{OR}^{21}$, $\text{S}(=\text{O})_2\text{R}^{21}$ and $\text{S}(=\text{O})_2\text{OR}^{21}$; and

L_3 is a leaving group; and all other symbols are as defined above;

in the presence of an organic or inorganic base in an organic solvent or a mixture of at least two different organic solvents, at a temperature ranging from -40°C to 150°C , for 0.5 to 16 h, to effect in situ cyclization to form the compound of general formula (I), wherein R^G is phenyl having at least one substituent which is $\text{OCH}_2\text{Phenyl}$, R^A is COOEt and s is 2; converting the $\text{OCH}_2\text{Phenyl}$ into hydroxyl and subsequently coupling the hydroxyl with the group $\text{L}_4(\text{CH}_2)_q\text{CR}^{23}\text{R}^{24}\text{COR}^{25}$, where L_4 is a leaving group;

optionally converting one or both of the COOEt groups into the cyano group $(\text{CH}_2)_p\text{CN}$, wherein p is as defined; optionally, subsequently converting at least one of the cyano groups into a group of the formula (3), as defined; and, optionally, converting the resultant compound into a physiologically tolerable salt.


Applicant: Bansi Lal et al.
Application No.: 10/574,982

33. (Previously presented) A pharmaceutical composition, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
34. (Previously presented) A pharmaceutical composition for inhibiting the binding of fibrinogen to blood platelets, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
35. (Previously presented) A pharmaceutical composition for inhibiting the binding of fibrinogen to blood platelets, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, in combination with an antithrombotic agent and a pharmaceutically acceptable carrier.
- 36 - 44. (Canceled)

Applicant: Bansi Lal et al.
Application No.: 10/574,982

Clean Claim Set


25. A compound of the general formula (I):



wherein

ring A is phenyl;

R^A is a group of formula (3):



wherein p is 0;

s is 1;

R¹ is selected from: H, hydroxy, alkyl, partially or fully fluorinated alkyl, alkoxy, alkenyl, alkynyl, carboxy, -C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle;

R³ and R⁴ are independently selected from: H, alkyl, partially or fully fluorinated alkyl, alkenyl, alkynyl, -C(=O)OR⁵, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle, -OR⁵, -SR⁵, -NR⁵R⁶, -S(=O)₂NR⁵R⁶, -S(=O)₂R⁵, -C(=O)R⁵, -C(=O)NR⁵R⁶, -C(=O)OR⁵, -C(=O)SR⁵, -OC(=O)R⁵, -OC(=O)OR⁵, -OC(=O)NR⁵R⁶, -OS(=O)₂R⁵, -S(C=O)NR⁵ and -OS(=O)₂NR⁵R⁶, or R³ and R¹ or R⁴, together with

Applicant: Bansi Lal et al.
Application No.: 10/574,982

the respective nitrogen atoms to which they are attached, form an unsubstituted or substituted 5-, 6- or 7- membered partially saturated or aromatic heterocycle, optionally having one or more additional heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and $-C(=O)OR^5$;

R^5 and R^6 are independently selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein each of said alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkylalkyl group optionally contains at least one hetero atom selected from: N, S and O anywhere in the chain, including the terminal position;

R^B is H;

Y^1 and Y^2 , together, are selected from: $=O$ and $=S$;

Z is N;

W is CH;

R^C is H;

n is 0, 1, 2 or 3;

R^D and R^E are independently selected from: H and an unsubstituted or substituted group selected from: alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkenyl, alkynyl, oxo, carboxy, $-C(=O)OR^5$, $-OR^{17}$, $-SR^{17}$, $-NR^{17}R^{18}$, $-NHC(=O)R^{17}$, $-NHC(=O)OR^{17}$, $-OC(=O)R^{17}$, $-SC(=O)R^{17}$, $-OS(=O)_2R^{17}$ and $-NHS(=O)_2R^{17}$;

R^{17} and R^{18} have the same meaning as R^5 and R^6 , defined above;

R^F is selected from: O, S and $N(OR^{19})$;

R^{19} has the same meaning as R^5 , defined above;

R^G is selected from: aryl, heteroaryl, and partially or fully saturated heterocycle,

Applicant: Bansilal et al.
Application No.: 10/574,982

where said aryl, heteroaryl and heterocycle are substituted by one or more groups of the formula (5):



and optionally, further substituted by one or more groups selected from: $-R^5$, halogen, $-CN$, $-SCN$, $-CNO$, $-OR^{21}$, $-OC(=O)R^{21}$, $-OS(=O)_2R^{21}$, $-OS(=O)_2NR^{21}R^{22}$, $-OC(=O)OR^{21}$, $-OC(=O)SR^{21}$, $-OC(=O)NR^{21}R^{22}$, $-SR^{21}$, $-S(=O)R^{21}$, $-NO_2$, $-NR^{21}(OR^{22})$, $-NR^{21}R^{22}$, $-NR^{21}C(=O)R^{22}$, $-N(R^{21})C(=O)OR^{22}$, $-N[S(=O)_2R^{21}]R^{23}$, $C(=O)OR^{21}$, $-S(=O)_2R^{21}$ and $-S(=O)_2OR^{21}$;

R^{21} has the same meaning as R^1 , defined above, and R^2 is selected from: H, hydroxy, alkyl, partially or fully fluorinated alkyl, alkoxy, alkenyl, alkynyl, carboxy, $-C(=O)OR^5$, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl and heterocycle:

T is selected from: $-CH_2$, O, S and NH;

q is 0, 1, 2 or 3;

R^{23} and R^{24} are independently selected from: H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocycle and $C(=O)R^{25}$, wherein said alkyl and alkenyl optionally contain at least one hetero atom selected from: O, S and N, in any position of the alkyl or alkenyl chain, and said alkyl and alkenyl are unsubstituted or substituted with at least one group selected from: $-OR^1$, $-OC(=O)R^1$, $-OS(=O)_2R^1$, $-S(=O)_2NR^1R^2$, $-OC(=O)OR^1$, $-OC(=O)SR^1$, $-OC(=O)NR^1R^2$, $-SR^1$, $-S(=O)R^1$, $-SC(=O)H$, $-SC(=O)OR^1$, $-NR^1(OR^2)$, $-NR^1R^2$, $-NR^1C(=O)R^2$, $-N(R^1)C(=O)OR^2$, $-NR^1S(=O)_2R^2$, $C(=O)OR^1$, $-S(=O)_2R^1$ and $-S(=O)_2OR^1$;

R^{25} is selected from: OR^5 , SR^5 , $-OCR^3R^4$ and $-NR^5R^6$, wherein R^3 , R^4 , R^5 and R^6 are as defined above and wherein optionally, R^3 and R^4 , together with the carbon to which they are attached, form an unsubstituted or substituted 5-, 6- or 7-membered saturated, partially saturated or aromatic heterocycle having one or

Applicant: Bansi Lal et al.
Application No.: 10/574,982

more heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and $-C(=O)OR^5$; and the group NR^5R^6 is, optionally, a heterocycle containing at least one additional heteroatom selected from: O, S, and N; in all its stereoisomeric and tautomeric forms and mixtures thereof in all ratios, and its pharmaceutically acceptable salts.

26. A compound according to claim 25, wherein

R^G is selected from: phenyl, piperidinyl and piperazinyl, and said phenyl, piperidinyl and piperazinyl are substituted by one or more groups of the formula (5):



and optionally, further substituted by one or more groups selected from: $-R^5$, halogen, $-CN$, $-SCN$, $-CNO$, $-OR^{21}$, $-OC(=O)R^{21}$, $-OS(=O)_2R^{21}$, $-OS(=O)_2NR^{21}R^{22}$, $-OC(=O)OR^{21}$, $-OC(=O)SR^{21}$, $-OC(=O)NR^{21}R^{22}$, $-SR^{21}$, $-S(=O)R^{21}$, $-NO_2$, $-NR^{21}(OR^{22})$, $-NR^{21}R^{22}$, $-NR^{21}C(=O)R^{22}$, $-N(R^{21})C(=O)OR^{22}$, $-N[S(=O)_2R^{21}]R^{23}$, $C(=O)OR^{21}$, $-S(=O)_2R^{21}$ and $-S(=O)_2OR^{21}$; and R^{21} and R^{22} are as defined in claim 25.

27. A compound according to claim 25, wherein

R_1 is hydrogen;

R_3 and R_4 are independently selected from: H, OH, $-C(O)OH$ and $-C(O)Oalkyl$;

$R^B = R^C = R^D = R^E =$ hydrogen;

Y^1 and Y^2 , together are $=O$;

n is the integer 0 or 1;

Applicant: Bansil Lal et al.
Application No.: 10/574,982

R^C is phenyl, substituted with one or more of the group of formula (5): T-(CH₂)_q-CR²³R²⁴-COR²⁵, wherein R²³ is H and R²⁴ is H, and, optionally, the compound is further substituted with one or more of the groups selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy, -C(=O)OR⁵, SR²¹, S(=O)₂R²¹ and -N(R²¹)-C(O)CH₃, -CH₂C(O)R²⁵;

and R²⁵ is selected from: OR⁵, OCR³R⁴ and NR⁵R⁶, wherein R³ and R⁴, together with the carbon to which they are attached form an unsubstituted or substituted 5-, 6- or 7- membered saturated, partially saturated or aromatic heterocycle having one or more heteroatoms selected from: N, O and S, wherein the substituents are selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy, -C(=O)OR⁵; and

R⁵, R⁶ and R²¹ are independently selected from: H, alkyl and phenyl.

28. A compound according to claim 25, wherein

R₁ is hydrogen;

R₃ and R₄ are independently selected from: H, OH, -C(O)OH and -C(O)Oalkyl;

R^B = R^C = R^D = R^E = hydrogen;

Y¹ and Y², together are =O;

n is the integer 0 or 1;

R^G is selected from: piperidinyl and piperazinyl, wherein said piperidinyl and piperazinyl are substituted with one or more of the group of formula (5): T-(CH₂)_q-CR²³R²⁴-COR²⁵, wherein R²³ is H and R²⁴ is H and, optionally, further substituted with one or more groups selected from: hydroxy, halogen, alkyl, alkoxy, alkenyl, alkynyl, oxo, carboxy and -C(=O)OR⁵;

and

R²⁵ is OR⁵, wherein R⁵ is selected from: H, alkyl and phenyl.

Applicant: Bansil Lal et al.
Application No.: 10/574,982

29. A compound according to claim 25 selected from:

- (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
- (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
- (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
- (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
- 4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
- (4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
- (4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
- (4-{2-[5-(Benzyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
- (4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
- (4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
- (4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
- (4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;

Applicant: Bansil Lal et al.
Application No.: 10/574,982

(4-{2-[5-(Benzyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methanesulfonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-ethoxy carbonyl methoxy-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(imino-{3-methyl-butyrylamino}-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-1-hydroxyimino-ethyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isobutoxy carbonyl methoxy-phenoxy)-acetic acid isobutyl ester;
2-(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-NN-diethyl-acetamide;

Applicant: Bansil Lal et al.
Application No.: 10/574,982

4-(2-{4-[2-(5-Carbamimidoyl-1-oxo-1,3-dihydro-isindol-2-yl)-acetyl]-phenoxy}-acetoxy)-piperidine-1-carboxylic acid benzyl ester;
4-Benzyloxycarbonylamino-2-(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-butyric acid ethyl ester;
4-Benzyloxycarbonylamino-2-(4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-butyric acid ethyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl}-phenylsulfanyl)-acetic acid methyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl}-2-chloro-phenoxy)-acetic acid ethyl ester;
(2-Chloro-4-{2-[5-(imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-acetic acid ethyl ester;
(2-Chloro-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl}-2-ethyl sulfanyl-phenoxy)-acetic acid ethyl ester;
(2-Ethylsulfanyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl}-2-ethane sulfonfyl-phenoxy)-acetic acid ethyl ester;
(2-Ethanesulfonfyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-acetic acid ethyl ester;
(2,6-Bis-ethylsulfanyl-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy}-acetic acid ethyl ester;
(2-Acetylamino-4-{2-[5-N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(imino-isobutoxy carbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-{2-[5-(N-hydroxy carbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;

(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid;

(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-methoxy-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-propoxy-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy carbonylmethoxy-phenoxy)-acetic acid ethyl ester;

(3-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid;

(2-Ethylsulfanyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-Ethyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(5-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isopropyl-phenoxy)-acetic acid ethyl ester;
(2-*tert*-Butyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid benzyl ester;
(2-Ethyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid;
(4-Hydroxy-3-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-5-methoxy-phenoxy)-acetic acid ethyl ester;
(3,5-Dihydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

Applicant: Bansil Lal et al.
Application No.: 10/574,982

(2-Ethoxycarbonylmethoxy-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(1-{2S-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-3-(4-hydroxy-phenyl)-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{3-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(5-Methyl-isoxazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(*tert*-Butoxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid;
(3-Ethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy-phenoxy)-acetic acid;
(3-Hydroxy-4-{2-[1-oxo-5-(5-oxo-2,5-dihydro-[1,2,4]oxadiazol-3-yl)-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(4-{2-[5-(Acetylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Acetoxy-4-{2-[5-(5-methyl-[1,2,4]oxadiazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid; and
(3-Allyloxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester.

30. A compound according to claim 27 selected from:

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid methyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(4-{2-[5-(Benzyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isopropyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Benzyloxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(Imino-methanesulfonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid isobutyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid;
(4-{2-[5-(Imino-methoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-(Imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-ethoxy carbonyl methoxy-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-Ethoxycarbonylmethoxy-4-{2-[5-(imino-{3-methyl-butyrylamino}-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-1-hydroxyimino-ethyl}-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isobutoxy carbonyl methoxy-phenoxy)-acetic acid isobutyl ester;

2-(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-NN-diethyl-acetamide;

4-(2-{4-[2-(5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl)-acetyl]-phenoxy}-acetoxy)-piperidine-1-carboxylic acid benzyl ester;

4-Benzyloxycarbonylamino-2-(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-butyric acid ethyl ester;

4-Benzyloxycarbonylamino-2-(4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-butyric acid ethyl ester;

(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenylsulfanyl)-acetic acid methyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-chloro-phenoxy)-acetic acid ethyl ester;

(2-Chloro-4-{2-[5-(imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(2-Chloro-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-ethyl sulfanyl-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(2-Ethylsulfanyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-2-ethanesulfonyl-phenoxy)-acetic acid ethyl ester;
(2-Ethanesulfonyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2,6-Bis-ethylsulfanyl-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2-Acetyl-amino-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-[2-[5-(imino-isobutoxycarbonylamino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(2-(Ethoxycarbonylmethyl-methanesulfonyl-amino)-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-[2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-phenoxy)-acetic acid benzyl ester;
(4-[2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-3-hydroxy-phenoxy)-acetic acid;
(4-[2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl]-3-methoxy-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(4-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-propoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy carbonylmethoxy-phenoxy)-acetic acid ethyl ester;
(3-Ethoxycarbonylmethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid;
(2-Ethylsulfanyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(5-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-isopropyl-phenoxy)-acetic acid ethyl ester;
(2-*tert*-Butyl-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Chloro-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-methyl-phenoxy)-acetic acid benzyl ester;
(2-Ethyl-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid ethyl ester;

Applicant: Bansi Lal et al.
Application No.: 10/574,982

(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid benzyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid;
(4-Hydroxy-3-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-5-methoxy-phenoxy)-acetic acid ethyl ester;
(3,5-Dihydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-3-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(2-Ethoxycarbonylmethoxy-5-hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(3-Ethoxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-ethoxy-phenoxy)-acetic acid;
(3-Hydroxy-4-{2-[1-oxo-5-(5-oxo-2,5-dihydro-[1,2,4]oxadiazol-3-yl)-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;
(4-{2-[5-(Acetyl)amino-imino-methyl]-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-phenoxy)-acetic acid ethyl ester;
(3-Acetoxy-4-{2-[5-(5-methyl-[1,2,4]oxadiazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester;


Applicant: Bansil Lal et al.
Application No.: 10/574,982

(4-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-3-hydroxy-2-propyl-phenoxy)-acetic acid ethyl ester;
(3-Hydroxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-2-propyl-phenoxy)-acetic acid; and
(3-Allyloxy-4-{2-[5-(N-hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-phenoxy)-acetic acid ethyl ester.


31. (Previously presented) A compound according to claim 28 selected from:
(1-{2S-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-3-(4-hydroxy-phenyl)-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{3-[5-(N-Hydroxycarbamimidoyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-propionyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(5-Methyl-isoxazol-3-yl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester;
(1-{2-[5-(*tert*-Butoxycarbonylamino-imino-methyl)-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid ethyl ester; and
(1-{2-[5-Carbamimidoyl-1-oxo-1,3-dihydro-isoindol-2-yl]-acetyl}-piperidin-4-yloxy)-acetic acid.

Applicant: Bansi Lal et al.
Application No.: 10/574,982

32. A process for the preparation of the compound of claim 25 having the general formula (I):




wherein all symbols have the same meaning as defined in claim 25, the process comprising: (a) reacting a compound of formula (II):



wherein

L is a leaving group; and all other symbols are as defined in claim 25; with a compound of the formula (III):




wherein all symbols are as defined in claim 25;

in the presence of an organic or inorganic base in an organic solvent or a mixture of at least two different organic solvents, at a temperature ranging from -40°C to 150°C , for 0.5 to 16 h, to effect in situ cyclization to form a compound of the general formula (I) above, and, optionally, converting the compound into a physiologically tolerable salt; or


Applicant: Bansi Lal et al.
Application No.: 10/574,982

b) reacting a compound of the formula (IV)



wherein

L₂ is a leaving group; and all other symbols are as defined in claim 25; with a compound of the formula (V):



where R^G is selected from: piperidinyl, piperazinyl and phenyl, wherein said piperidinyl, piperazinyl and phenyl, are optionally substituted with 1, 2, 3 or 4 hydroxyl groups, and all other symbols are as defined in claim 25, in the presence of an organic or inorganic base in an organic solvent or water at a temperature ranging from 0°C to 150°C, for 0.5 to 12 h, to form a compound of the general formula (I), and, optionally, converting one or more of the hydroxyl groups into a group selected from the substituents for R^G as defined in general formula (I) and, optionally, converting the compound into a physiologically tolerable salt; alternatively, activating a compound of the formula (IV) above, wherein L₂ is -OH, by treatment with a mixed anhydride to form a peptide coupling with a compound of the formula (V), wherein R^G is piperidinyl or piperazinyl, and thereby provide a compound of the general formula (I), wherein R^G is piperidinyl or piperazinyl substituted with at least a group of the formula (5); and, optionally, converting the resultant compound into a physiologically tolerable salt.

Applicant: Bansi Lal et al.
Application No.: 10/574,982

33. A pharmaceutical composition, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
34. A pharmaceutical composition for inhibiting the binding of fibrinogen to blood platelets, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
35. A pharmaceutical composition for inhibiting the binding of fibrinogen to blood platelets, comprising a compound of formula (I) according to claim 25, or a pharmaceutically acceptable salt thereof, in combination with an antithrombotic agent and a pharmaceutically acceptable carrier.

Applicant: Bansil Lal et al.
Application No.: 10/574,982

Conclusion:

Applicant believes that the foregoing proposed amendments would place the application in condition for allowance. Applicant requests entry of the amendments and rejoinder of claim 32. If any further matters need to be addressed, please contact Applicant's representative.

Respectfully submitted,

Bansil Lal et al.

By /Douglas J. Bucklin/
Douglas J. Bucklin
Registration No. 51,208

Volpe and Koenig, P.C.
United Plaza, Suite 1600
30 South 17th Street
Philadelphia, PA 19103
Telephone: (215) 568-6400
Facsimile: (215) 568-6499

DJB/dmp